



The effects of dietary cholecalciferol and 1α -hydroxycholecalciferol levels in a calcium- and phosphorus-deficient diet on growth performance and tibia quality of growing broilers

J.C. Han^{1,6}, H.X. Qu¹, J.Q. Wang², J.H. Yao³, C.M. Zhang¹, G.L. Yang¹, Y.H. Cheng⁴ and X.S. Dong⁵

¹Shangqiu Normal University, Institute of Animal Nutrition and Animal Genetics, College of Life Science, Shangqiu 476000, P.R. China

²Shenyang Agricultural University, College of Science and Technology, Shenyang 113122, P.R. China

³Northwest A&F University, College of Animal Science and Technology, Yangling 712100, P.R. China

⁴I-Lan University, Department of Biotechnology and Animal Science, I-Lan 26047, Taiwan, ROC

⁵Taizhou Healtch Chemical Co., Ltd., Taizhou 318000, P.R. China

KEY WORDS: 1α -hydroxycholecalciferol, cholecalciferol, growth, tibia, potency, broiler

ABSTRACT. This study aims to evaluate the effects of cholecalciferol (vitamin D_3 , 0, 5, 10, 25, 50, 100, 250, 500, and 1000 $\mu\text{g} \cdot \text{kg}^{-1}$) and 1α -hydroxycholecalciferol (1α -OH D_3 , 0, 2.5, and 5 $\mu\text{g} \cdot \text{kg}^{-1}$) on growth performance and tibia quality and to compare the relative potency of 1α -OH D_3 to vitamin D_3 in 1- to 21-day-old female broilers fed a calcium (Ca)- and phosphorus (P)-deficient diet. The basal diet contained 0.50% Ca, 0.25% non-phytate phosphorus (NPP) and was not supplemented with vitamin D_3 , whereas the control diet contained 1.00% Ca, 0.45% NPP, and 25 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D_3 . Dietary vitamin D_3 levels significantly affected body weight gain (BWG), feed intake (FI), serum Ca and tibia parameters. The addition of 5 $\mu\text{g} \cdot \text{kg}^{-1}$ 1α -OH D_3 resulted in greater BWG, FI, and serum Ca as well as tibia ash weight and content, breaking strength, and Ca and P contents, compared with the birds fed the basal diet. Using BWG, serum Ca, tibia ash weight and content, and tibia Ca and P content as the criteria, the vitamin D_3 requirements of 1- to 21-day-old broilers fed Ca- and P-deficient diets were 64.0, 16.2, 173.0, 65.1, 33.1 and 30.0 $\mu\text{g} \cdot \text{kg}^{-1}$, and the relative potencies of 1α -OH D_3 to vitamin D_3 were 5.03, 2.19, 18.00, 5.14, 4.09 and 3.33, respectively. These data indicate that high levels of vitamin D_3 can spare the use of P in broiler diets and that the potency of 1α -OH D_3 is higher than that of vitamin D_3 .

Received: 29 February 2012

Revised: 23 April 2013

Accepted: 21 June 2013

⁶ Corresponding author: e-mail: j.c.han@hotmail.com

Introduction

In a basal diet with low levels of calcium (Ca, 0.64%) and total phosphorus (tP, 0.42%), 8- to 20-day-old broilers that were fed 1250 $\mu\text{g} \cdot \text{kg}^{-1}$ of cholecalciferol (vitamin D_3) achieved greater body weight gain (BWG) and tibia ash content than birds

fed 37.5 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D_3 (Baker et al., 1998). On the other hand, chickens fed Ca- and P-adequate diets (1.00% Ca, 0.77% tP) with 5 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D_3 exhibited BWG and bone ash values as high as those of chickens that received 30 and 1250 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D_3 (Baker et al., 1998). These results indicate that broilers require less vitamin D_3

when their diet contains adequate Ca and P. When the broilers were fed sufficient Ca and P, their body weight and tibia breaking strength were maximized on day 14 with $250 \mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D_3 , tibia ash was maximized with $125 \mu\text{g} \cdot \text{kg}^{-1}$ vitamin D_3 (Whitehead et al., 2004), and bone area, thickness, and mineral content increased with $250 \mu\text{g} \cdot \text{kg}^{-1}$ vitamin D_3 (Kim et al., 2011). With growth improvement, broilers need more vitamin D for bone development and strength. In addition, the environmental pollution caused by P in poultry production requires the increasing attention of animal nutritionists. Qian et al. (1997) found that dietary Ca to tP ratios ranging from 1.1 to 1.4 are critical for the efficient use of phytase and vitamin D_3 in improving the utilization of phytate P. Rao et al. (2006) reported that in a basal diet with 0.50% Ca and 0.25% of non-phytate phosphorus (NPP), the addition of $90 \mu\text{g} \cdot \text{kg}^{-1}$ vitamin D_3 maintains growth performance and bone mineralization in broilers. These data suggest that the addition of high levels of vitamin D_3 can spare the use of P in broiler diets with an appropriate Ca to P ratio. The vitamin D_3 requirement of broilers for growth and bone quality in Ca- and P-sub-adequate diets is unclear, however.

The compounds 1α -hydroxycholecalciferol (1α -OH D_3) and 25-hydroxycholecalciferol (25-OH D_3) are vitamin D metabolites. Recently, 25-OH D_3 has been approved for use in feed for fattening chickens, laying hens and turkeys in Europe. It is possible that 1α -OH D_3 will become a feed additive in the future. Kasim and Edwards (2000) evaluated the effects of different vitamin D_3 sources on growth and tibia ash using broiler chickens fed Ca- and P-adequate diets (1.0% Ca, 0.7% tP). In other studies, similar basal diets with sufficient Ca and P were used to compare the efficacy of vitamin D_3 with that of 1α -OH D_3 (Edwards et al., 2002) and that of vitamin D_3 with that of 25-OH D_3 (Yarger et al., 1995; Fritts and Waldroup, 2003). Vitamin D_3 (Qian et al., 1997) and 1α -OH D_3 (Han et al., 2012) have high activity at low levels of dietary Ca and P. Thus, comparing the efficacy of vitamin D metabolites should be conducted under Ca- and P-deficient diets. In the present study, the effect of vitamin D_3 and 1α -OH D_3 on growth performance and tibia quality was reevaluated on the basis of diets with 0.50% Ca and 0.25% NPP.

The objective of the presented study is to determine the vitamin D_3 requirement of 1- to 21-day-old broilers fed Ca- and P-deficient diets and to reevaluate the relative potency of 1α -OH D_3 compared with that of vitamin D_3 .

Material and methods

Birds, diets, management

All procedures used in the present experiment were approved by the Animal Care Committee of Shangqiu Normal University. On the day of hatching, 480 female Ross broilers were individually weighed and randomly assigned to 12 treatments, four cages (70 cm \times 70 cm \times 30 cm) with 10 birds per treatment. The lighting system consisted of 23 h of light from days 0 to 21. The room temperature was maintained at 33°C from days 0 to 3 and then gradually reduced by 3°C per week to a final temperature of 24°C.

Twelve experimental diets were prepared. Vitamin D_3 and 1α -OH D_3 were supplied by Tongxiang Tianhecheng Food Technology Co., Ltd. (Jiaxing, China) and Taizhou Healtech Chemical Co., Ltd. (Taizhou, China), respectively. The crystalline vitamin D_3 and 1α -OH D_3 were dissolved in ethanol and then brought to final concentrations of $200 \text{ mg} \cdot \text{l}^{-1}$ of vitamin D_3 and $10 \text{ mg} \cdot \text{l}^{-1}$ of 1α -OH D_3 in a solution of 5% ethanol and 95% propylene glycol (Han et al., 2009).

The basal diet contained 0.50% Ca, 0.25% NPP and was without added vitamin D_3 . It was supplemented with 5, 10, 25, 50, 100, 250, 500 and 1000 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D_3 or 2.5 and $5 \mu\text{g} \cdot \text{kg}^{-1}$ of 1α -OH D_3 (Table 1). A diet with adequate

Table 1. Ingredients and nutrient composition of the experimental diets

Component	Diet ³	
	basal	control
Ingredient, g \cdot kg ⁻¹		
maize	638.7	600.9
soya bean meal	290.0	290.0
soya bean oil	10.0	10.0
lard	7.2	20.5
soya bean protein isolate	31.3	36.5
limestone	6.9	13.9
dicalcium phosphate	7.2	19.5
L-lysine-HCl	1.8	1.8
DL-methionine	0.6	0.6
trace mineral premix ¹	1.0	1.0
vitamin premix ²	0.3	0.3
choline chloride	2.0	2.0
NaCl	3.0	3.0
Nutrient composition		
metabolizable energy, MJ \cdot kg ⁻¹	12.55	12.55
analysed crude protein, %	20.5	20.4
analysed calcium, %	0.55	1.08
analysed total phosphorus, %	0.50	0.68
calculated non-phytate phosphorus, %	0.25	0.45

¹ the trace-mineral premix provided the following, per kg of diet: mg: iron 100, zinc 100, copper 8, manganese 120, iodine 0.7, selenium 0.3; ² the vitamin premix provided the following, per kg of diet: IU: vit. A 8000, vit. E 20; mg: menadione 0.5, thiamine 2.0, riboflavin 8.0, niacin 3.5, pyridoxine 0.01; vit. B₁₂ 0.01, pantothenic acid 10.0, folic acid 0.55, biotin 0.18; ³ to the control diet $25 \mu\text{g} \cdot \text{kg}^{-1}$ vit. D_3 was added

levels of Ca (1.00%), NPP (0.45%) and vitamin D₃ (25 µg · kg⁻¹) was used as the control. The birds were given access to mash diets and water *ad libitum* during the 21-day experiment.

Sample collection

On day 21, the broilers were individually weighed. Two chickens were randomly chosen from each replicate for blood and tibia collection. Serum samples (5 ml) were collected using 5 ml anticoagulated syringes (Shanghai K&G International Co. Ltd., Shanghai, China) by cardiac puncture and then centrifuged for 10 min at 3000 g at 20°C.

The birds were sacrificed by exsanguination, plucked, eviscerated, abdominal fat was removed, and the carcasses were weighed. The left and right tibias were excised and frozen at -20°C for further analysis (breaking strength, ash weight, and content of ash, Ca, and P).

Sample analysis

The Ca and inorganic phosphate (Pi) in blood serum were immediately determined using a Shimadzu CL-8000 Analyzer (Shimadzu Corp., Kyoto, Japan) following the manufacturer's instructions. According to the method by Hall et al. (2003), the left tibia was boiled for 5 min to loosen the muscle tissue. The meat, connective tissue, and fibula bone were removed from the tibia. The tibia was placed in a container with ethanol for 48 h to remove water and polar lipids. The bones were then further extracted in anhydrous ether for 48 h to remove non-polar lipids. The tibias were dried at 105°C for 24 h, weighed, and their weight recorded. Tibia ash content was determined by ashing the bone in a muffle furnace for 18 h at 600°C.

The right tibia was used to analyse the breaking strength. The tibia breaking strength was determined using an all-digital electronic universal testing machine (Shenzhen Hengen Instrument Co. Ltd., Shenzhen, China). The tibias were cradled on two support points 4 cm apart. Using a 50 kg load cell and a crosshead speed of 10 mm · min⁻¹, the force was applied to the midpoint of the same facet of each tibia (Jendral et al., 2008).

Ca in diets was determined according to the EDTA titration method. Total P in diets was determined by photometric methods after reaction with ammonium molybdate and ammonium metavanadate. The crude protein content in diets was determined according to the Kjeldhal method on a PN-1430 apparatus (Barcelona, Spain).

Statistical analyses

Replicate means are the experimental units in the statistical analysis. All data were analysed with an ANOVA procedure of SAS software (SAS Institute, 2001). The means were compared with Tukey's test at $p < 0.05$. Nonlinear regression analysis was conducted using the PROC NLIN procedure of SAS to estimate the vitamin D₃ requirement based on the selected variables, including BWG, serum Ca, tibia ash weight and content, and tibia Ca and P contents. The model was: $y = ax^2 + bx + c$, where y is the response and x is the variable. The relative potencies of 1α-OH D₃ to vitamin D₃ were determined by the slope ratio method (Littell et al., 1997). The model is: $y = a + b_1x_1 + b_2x_2$, where y is the response, x_1 is vitamin D₃, and x_2 is 1α-OH D₃.

Results

The levels of dietary vitamin D₃ and 1α-OH D₃ significantly affected BWG, feed intake (FI), and serum Ca but did not influence the feed conversion ratio (FCR), slaughter yield (SY), and serum Pi in 1- to 21-day-old broilers (Table 2). BWG and FI responded linearly to vitamin D₃ ranging from 0 to 50 µg · kg⁻¹ and to 1α-OH D₃ from 0 to 5 µg · kg⁻¹. Serum Ca also increased linearly to vitamin D₃ ranging from 0 to 10 µg · kg⁻¹ and to 1α-OH D₃ from 0 to 5 µg · kg⁻¹. Serum Pi in broiler chickens fed 25 µg · kg⁻¹ of vitamin D₃ was higher than that of birds fed the basal diet ($p < 0.05$). No significant differences were observed in BWG, FI, or serum Ca among the broilers fed 10 to 1000 µg · kg⁻¹ vitamin D₃ or 5 µg · kg⁻¹ 1α-OH D₃, and those fed the control diets.

The tibia ash weight of the broilers increased linearly from 0.42 to 0.71 g/tibia when the dietary vitamin D₃ increased from 0 to 100 µg · kg⁻¹ (Table 3). Tibia ash content (g · 100 g⁻¹) also enhanced linearly to vitamin D₃ ranging from 0 to 25 µg · kg⁻¹. Broilers fed 25 to 1000 µg · kg⁻¹ vitamin D₃ achieved a significantly greater tibia breaking strength than the birds fed the basal diet. The content of tibia Ca and P in broilers fed 25 µg · kg⁻¹ vitamin D₃ was significantly higher than that of birds fed the basal diet ($p < 0.05$). The addition of 5 µg · kg⁻¹ 1α-OH D₃ resulted in greater tibia ash weight and content, breaking strength, and Ca and P content, compared with the birds fed the basal diet. No significant differences in tibia ash weight, breaking strength, or Ca and P contents were observed between the broilers fed with 5 µg · kg⁻¹ 1α-OH D₃ or 25 to 1000 µg · kg⁻¹ vitamin D₃ and the control diets.

Using BWG, serum Ca, tibia ash weight and content, and tibia Ca and P contents as the criteria,

Table 2. Effects of vitamin D₃ and 1 α -OH D₃ on growth performance of 1- to 21-day-old broilers

Group	D ₃ $\mu\text{g} \cdot \text{kg}^{-1}$	1 α -OH D ₃ $\mu\text{g} \cdot \text{kg}^{-1}$	Growth ²				Serum ³		
			BWG g · bird ⁻¹	FI g · bird ⁻¹	FCR (FI/BWG)	SY %	Ca mg · 100 ml ⁻¹	Pi mg · 100 ml ⁻¹	
1	Basal diet		563 ^d	910 ^c	1.62	68.89	6.50 ^c	5.47 ^b	
2	5		581 ^{cd}	938 ^{bc}	1.62	68.05	7.55 ^b	5.83 ^{ab}	
3	10		617 ^{abc}	1041 ^{ab}	1.69	70.05	8.64 ^a	6.11 ^{ab}	
4	25		638 ^{ab}	1070 ^a	1.68	69.79	8.14 ^{ab}	6.75 ^a	
5	50		641 ^{ab}	1082 ^a	1.69	69.10	8.20 ^{ab}	6.25 ^{ab}	
6	100		634 ^{ab}	1052 ^a	1.66	69.07	8.54 ^{ab}	6.41 ^{ab}	
7	250		635 ^{ab}	1099 ^a	1.73	68.07	8.27 ^{ab}	6.21 ^{ab}	
8	500		638 ^{ab}	1073 ^a	1.68	68.95	8.53 ^{ab}	6.45 ^{ab}	
9	1000		647 ^{ab}	1085 ^a	1.68	68.67	8.41 ^{ab}	6.38 ^{ab}	
10		2.5	602 ^{bcd}	1023 ^{abc}	1.70	68.86	8.36 ^{ab}	5.90 ^{ab}	
11		5.0	618 ^{abc}	1041 ^{ab}	1.69	70.40	8.45 ^{ab}	5.82 ^{ab}	
12	Control diet		648 ^a	1067 ^a	1.65	71.83	8.69 ^a	6.72 ^a	
SEM			4	10	0.01	0.25	0.10	0.08	
P value			<0.001	<0.001	0.435	0.092	<0.001	0.012	
Source of variation			Probabilities						
Vitamin D ₃ level			<0.001	<0.001	0.392	0.669	<0.001	0.057	
1 α -OH D ₃ level			0.021	0.006	0.313	0.368	<0.001	0.400	

^{a-d} means in the same column without a common superscript differ significantly ($p < 0.05$); ¹ the basal diet contained 0.50% Ca and 0.25% NPP, no added vitamin D₃. The control diet contained 1.00% Ca, 0.45% NPP, and 25 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃; BWG – body weight gain, FI – feed intake, FCR – feed conversion ratio, SY – slaughter yield (carcass weight as a percentage of live body weight), Ca – calcium, Pi – inorganic phosphate; ² data are means of 4 replicate cages consisting of 10 birds per replicate cage; ³ data are means of 4 replicates consisting of 2 birds per replicate

Table 3. Effects of vitamin D₃ and 1 α -OH D₃ on tibia quality of 1- to 21-day-old broilers

Group	D ₃ $\mu\text{g} \cdot \text{kg}^{-1}$	1 α -OH D ₃ $\mu\text{g} \cdot \text{kg}^{-1}$	Ash		Breaking- strength, N	Ca g · 100 g ⁻¹	P g · 100 g ⁻¹		
			g/tibia	g · 100 g ⁻¹					
1	Basal diet		0.42 ^c	40.96 ^d	43.67 ^d	14.12 ^b	6.73 ^b		
2	5		0.55 ^{bc}	41.34 ^{cd}	63.48 ^{bcd}	15.13 ^{ab}	7.39 ^{ab}		
3	10		0.58 ^{bc}	42.48 ^{bcd}	61.13 ^{cd}	15.44 ^{ab}	7.70 ^a		
4	25		0.61 ^{ab}	44.05 ^{ab}	77.20 ^{abc}	16.19 ^a	8.06 ^a		
5	50		0.65 ^{ab}	42.65 ^{bcd}	69.45 ^{abc}	15.89 ^{ab}	7.62 ^{ab}		
6	100		0.71 ^{ab}	43.24 ^{bc}	78.30 ^{abc}	16.15 ^a	7.65 ^{ab}		
7	250		0.66 ^{ab}	43.61 ^{ab}	73.67 ^{abc}	15.16 ^{ab}	7.83 ^a		
8	500		0.62 ^{ab}	42.16 ^{bcd}	67.59 ^{abc}	15.21 ^{ab}	7.70 ^a		
9	1000		0.67 ^{ab}	43.61 ^{ab}	80.54 ^{abc}	15.14 ^{ab}	7.79 ^a		
10		2.5	0.67 ^{ab}	43.33 ^{bc}	69.55 ^{abc}	15.27 ^{ab}	7.54 ^{ab}		
11		5.0	0.69 ^{ab}	43.78 ^{ab}	86.84 ^a	15.98 ^a	7.72 ^a		
12	Control diet		0.77 ^a	45.57 ^a	82.51 ^{ab}	16.49 ^a	8.18 ^a		
SEM			0.02	0.21	1.95	0.13	0.07		
P value			<0.001	<0.001	<0.001	0.003	0.002		
Source of variation			Probabilities						
Vitamin D ₃ level			<0.001	<0.001	<0.001	0.003	0.008		
1 α -OH D ₃ level			<0.001	0.002	<0.001	0.087	0.057		

^{a-d} means in the same column without a common superscript differ significantly ($p < 0.05$); ¹ the basal diet contained 0.50% Ca and 0.25% NPP, no added vitamin D₃. The control diet contained 1.00% Ca, 0.45% NPP, and 25 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃; data are means of 4 replicates consisting of 2 birds per replicate; Ca – calcium, P – phosphorus

Table 4. Vitamin D₃ requirements of 1- to 21-day-old broilers fed Ca- and P-deficient diets

Indices	Formula	P value	X, D ₃ , $\mu\text{g} \cdot \text{kg}^{-1}$	D ₃ requirement, $\mu\text{g} \cdot \text{kg}^{-1}$
Body weight gain	$y = -0.02050x^2 + 2.6250x + 574.5$	<0.001	0 to 100	64.0
Serum calcium	$y = -0.00939x^2 + 0.3035x + 6.4335$	<0.001	0 to 25	16.2
Tibia ash weight	$y = -0.00001x^2 + 0.00346x + 0.5044$	<0.001	0 to 250	173.0
Tibia ash content	$y = -0.00056x^2 + 0.0729x + 41.3806$	0.002	0 to 100	65.1
Tibia calcium	$y = -0.00194x^2 + 0.1283x + 14.3039$	0.002	0 to 50	33.1
Tibia phosphorus	$y = -0.00149x^2 + 0.0893x + 6.8624$	0.003	0 to 50	30.0

Table 5. Relative potencies of 1 α -OH D₃ to vitamin D₃ in 1- to 21-day-old broilers fed Ca- and P-deficient diets¹

Indices	Formula	P value	x ₁ $\mu\text{g} \cdot \text{kg}^{-1}$	x ₂ $\mu\text{g} \cdot \text{kg}^{-1}$	Relative potency
Body weight gain	$y = 582.1757 + 1.4412x_1 + 7.2444x_2$	< 0.001	0 to 50	0 to 5	5.03
Serum calcium	$y = 6.7471 + 0.1835x_1 + 0.4015x_2$	< 0.001	0 to 10	0 to 5	2.19
Tibia ash weight	$y = 0.5247 + 0.0021x_1 + 0.0378x_2$	< 0.001	0 to 100	0 to 5	18.00
Tibia ash content	$y = 41.1253 + 0.1170x_1 + 0.6018x_2$	< 0.001	0 to 25	0 to 5	5.14
Tibia calcium	$y = 14.4813 + 0.0740x_1 + 0.3025x_2$	0.006	0 to 25	0 to 5	4.09
Tibia phosphorus	$y = 7.0305 + 0.0456x_1 + 0.1518x_2$	0.003	0 to 25	0 to 5	3.33

¹x₁ = D₃, x₂ = 1 α -OH D₃

the vitamin D₃ requirements of 1- to 21-day-old broilers fed Ca- and P-deficient diets were 64.0, 16.2, 173.0, 65.1, 33.1 and 30.0 $\mu\text{g} \cdot \text{kg}^{-1}$ (Table 4), and the relative potencies of 1 α -OH D₃ to vitamin D₃ were 5.03, 2.19, 18.00, 5.14, 4.09 and 3.33, respectively (Table 5).

Discussion

The addition of 60 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D₃ increased the BWG of broilers fed with Ca- and P-deficient diets, reaching levels similar to those of birds fed control diets (Rao et al., 2006). Driver et al. (2005) also found that Ca- and P-deficient broilers fed 1 α -OH D₃ and phytase had equal growth performance and tibia ash weight compared with birds fed diets with adequate Ca and P. In the present study, no significant difference was observed in the growth rate as well as ash weight, breaking strength, and Ca and P contents of the tibia among the broilers fed 25 to 1000 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃ and those fed the control diets.

Phosphorous deficiency reduces carcass weight (Angel et al., 2006) and yield in broilers (Chen and Moran, 1995). When phytase was added, the carcass yield increased (Pillai et al., 2006). By contrast, the slaughter yield of the broilers was not significantly affected by the supplementation of vitamin D₃ or 1 α -OH D₃ in the present experiment. Scheideler and Ferket (2000) also found that the dietary P levels (NPP 0.30 vs. 0.42%) did not influence the weight of the carcass, thigh and breast meat of the broilers.

Dietary vitamin D₃ (Rao et al., 2006), 1,25-(OH)₂ D₃ (Edwards, 2002), and 1 α -OH D₃ (Edwards et al., 2002; Han et al., 2009) increased the serum Ca and Pi concentrations in broilers. In contrast, Driver et al. (2008) reported that 1 α -OH D₃ did not affect the serum Pi content in non-obese diabetic mice. In the present study, the increase of dietary vitamin D₃ enhanced serum Ca and Pi contents, whereas serum P did not depend on the 1 α -OH D₃ level.

Studies have shown that vitamin D₃ (Rao et al., 2009), 1 α -OH D₃ (Snow et al., 2004),

25-OH D₃ (Aburto et al., 1998) and 1,25-(OH)₂ D₃ (Edwards, 2002) increase bone weight, length, and ash in broilers. These data indicate that vitamin D metabolites increase bone growth and mineral deposition in poultry. Fritts and Waldroup (2003) reported no significant differences in tibia ash weight among the broilers fed 25, 50 and 100 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃. Similar results were found in the present experiment. When the vitamin D₃ level ranged from 25 to 1000 $\mu\text{g} \cdot \text{kg}^{-1}$, no significant differences were observed in tibia ash weight, breaking strength, and Ca and P contents of the tibia of the broilers.

The NRC (1994) recommendation on vitamin D₃ for broilers is 5 $\mu\text{g} \cdot \text{kg}^{-1}$ in Ca- and P-adequate diets, which may be lower than the practical requirement. Research has shown that about 25 and 50 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D₃ were needed to maximize the body weight and bone ash of 42-day-old broilers, respectively (Fritts and Waldroup, 2003). When broilers were fed with Ca- and P-deficient diets, the addition of 1250 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃ resulted in greater tibia ash than the addition of 0 and 37.5 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D₃ (Baker et al., 1998). The addition of 6660 $\mu\text{g} \cdot \text{kg}^{-1}$ vitamin D₃ yielded higher toe ash contents and Ca and P retention than the addition of 66 and 660 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D₃ (Qian et al., 1997). At 42 d, the body weight of broilers fed 100 $\mu\text{g} \cdot \text{kg}^{-1}$ of vitamin D₃ was significantly greater than of birds fed 25 or 50 $\mu\text{g} \cdot \text{kg}^{-1}$ (Fritts and Waldroup, 2005). These data suggest that the vitamin D₃ requirements of broilers in the Ca- and P-deficient diets were more than those of the birds fed Ca- and P-adequate diets. Rao et al. (2009) reported, however, that the vitamin D₃ requirements of broilers fed Ca- and P-deficient diets ranged from 16 to 25 $\mu\text{g} \cdot \text{kg}^{-1}$. This value may be lower than the actual requirement. Using weight gain and tibia ash weight and content as the criteria in the present experiment, we found that the vitamin D₃ requirements were 64.0, 173.0 and 65.1 $\mu\text{g} \cdot \text{kg}^{-1}$ in broilers fed Ca- and P-deficient diets, respectively.

Previous research has shown that 1 α -OH D₃ is 4.5 times more active than vitamin D₃ in respect to the BWG of broiler chickens (Soares et al., 1978),

8.6 times in respect to tibia ash increase (Boris et al., 1977), and at least 10 times in terms of mobilizing bone calcium and raising plasma calcium concentrations in Leghorn cockerels (Haussler et al., 1973). Using body weight, plasma Ca, tibia ash weight and percentage, and incidence of rickets as criteria, Edwards et al. (2002) reported that the relative biological values of 1α -OH D_3 to vitamin D_3 were 10.08, 9.50, 11.26, 4.48, and 4.50, respectively. The relative potency of 1α -OH D_3 to vitamin D_3 on body weight in our study was similar to the results of Soares et al. (1978) and lower in BWG and plasma Ca, whereas higher in tibia ash weight and content, than those reported by Edwards et al. (2002). The differences may have been caused by the age and sex of the birds and the dietary Ca and P contents. Edwards et al. (2002) used 1- to 16-day-old mixed-sex broilers fed 1.0% Ca and 0.7% tP, whereas the present experiment used 1- to 21-day-old female birds fed 0.50% Ca and 0.48% tP. Our results are in accordance with other studies where the relative efficacy of 1α -OH D_3 to vitamin D_3 in terms of tibia ash weight was higher than for body weight.

Conclusions

Dietary vitamin D_3 and 1α -OH D_3 improved growth performance and serum Ca as well as the tibia quality of broiler chickens. Using BWG, serum Ca, tibia ash weight and content, and tibia Ca and P contents as the criteria, the vitamin D_3 requirements of 1- to 21-day-old broilers fed Ca- and P-deficient diets were 64.0, 16.2, 173.0, 65.1, 33.1 and 30.0 $\mu\text{g} \cdot \text{kg}^{-1}$, and the relative potencies of 1α -OH D_3 to vitamin D_3 were 5.03, 2.19, 18.00, 5.14, 4.09 and 3.33, respectively. These data indicate that high levels of vitamin D_3 can spare the use of P in broiler diets and that the potency of 1α -OH D_3 is higher than that of vitamin D_3 .

Acknowledgements

Supported by the National Natural Science Foundation of China (No. 31101732) and the Natural Science Foundation of Henan Province (No. 2011B230011).

References

- Aburto A., Jr Edwards H.M., Britton W.M., 1998. The influence of vitamin A on the utilization and amelioration of toxicity of cholecalciferol, 25-hydroxycholecalciferol, and 1,25-dihydroxycholecalciferol in young broiler chickens. *Poultry Sci.* 77, 585–593
- Angel R., Saylor W.W., Mitchell A.D., Powers W., Applegate T.J., 2006. Effect of dietary phosphorus, phytase, and 25-hydroxycholecalciferol on broiler chicken bone mineralization, litter phosphorus, and processing yields. *Poultry Sci.* 85, 1200–1211
- Baker D.H., Biehl R.R., Emmert J.L., 1998. Vitamin D_3 requirement of young chicks receiving diets varying in calcium and available phosphorus. *Brit. Poultry Sci.* 39, 413–417
- Boris A., Hurley J.F., Trmal T., 1977. Relative activities of some metabolites and analogs of cholecalciferol in stimulation of tibia ash weight in chicks otherwise deprived of vitamin D. *J. Nutr.* 107, 194–198
- Chen X., Jr. Moran E.T., 1995. The withdrawal feed of broilers: Carcass responses to dietary phosphorus. *J. Appl. Poultry Res.* 4, 69–82
- Driver J.P., Foreman O., Mathieu C., van Etten E., Serreze D.V., 2008. Comparative therapeutic effects of orally administered 1,25-dihydroxyvitamin D_3 and 1alpha-hydroxyvitamin D_3 on type-1 diabetes in non-obese diabetic mice fed a normal-calcaemic diet. *Clin. Exp. Immunol.* 151, 76–85
- Driver J.P., Pesti G.M., Bakalli R.I., Edwards H.M. Jr., 2005. Phytase and 1 α -hydroxycholecalciferol supplementation of broiler chickens during the starting and growing/finishing phases. *Poultry Sci.* 84, 1616–1628
- Edwards Jr. H.M., 2002. Studies on the efficacy of cholecalciferol and derivatives for stimulating phytate utilization in broilers. *Poultry Sci.* 81, 1026–1031
- Edwards Jr. H.M., Shirley R.B., Escoe W.B., Pesti G.M., 2002. Quantitative evaluation of 1 α -hydroxycholecalciferol as a cholecalciferol substitute for broilers. *Poultry Sci.* 81, 664–669
- Fritts C.A., Waldroup P.W., 2003. Effect of source and level of vitamin D on live performance and bone development in growing broilers. *J. Appl. Poultry Res.* 12, 45–52
- Fritts C.A., Waldroup P.W., 2005. Comparison of cholecalciferol and 25-hydroxycholecalciferol in broiler diets designed to minimize phosphorus excretion. *J. Appl. Poultry Res.* 14, 156–166
- Hall L.E., Shirley R.B., Bakalli R.I., Aggrey S.E., Pesti G.M., Edwards Jr. H.M., 2003. Power of two methods for the estimation of bone ash of broilers. *Poultry Sci.* 82, 414–418
- Han J.C., Liu Y., Yao J.H., Wang J.Q., Qu H.X., Yan Y.F., Yue J., Ding J.L., Shi Z.T., Dong X.S., 2012. Dietary calcium levels reduce the efficacy of one alpha-hydroxycholecalciferol in phosphorus-deficient diets of broilers. *J. Poultry Sci.* 49, 34–38
- Han J.C., Yang X.D., Zhang T., Li H., Li W.L., Zhang Z.Y., Yao J.H., 2009. Effects of 1 α -hydroxycholecalciferol on growth performance, parameters of tibia and plasma, meat quality, and type IIb sodium phosphate cotransporter gene expression of one- to twenty-one-day-old broilers. *Poultry Sci.* 88, 323–329
- Haussler M.R., Zerwekh J.E., Hesse R.H., Rizzardo E., Pechet M.M., 1973. Biological activity of 1 α -hydroxycholecalciferol, a synthetic analog of the hormonal form of vitamin D_3 . *Proc. Nat. Acad. Sci. USA.* 70, 2248–2252
- Jendral M.J., Korver D.R., Church J.S., Feddes J.J.R., 2008. Bone mineral density and breaking strength of white leghorns housed in conventional, modified, and commercially available colony battery cages. *Poultry Sci.* 87, 828–837
- Kasim A.B., Edwards Jr. H.M. Jr., 2000. Evaluation of cholecalciferol sources using broiler chick bioassays. *Poultry Sci.* 79, 1617–1622
- Kim W.K., Bloomfield S.A., Ricke S.C., 2011. Effects of age, vitamin D_3 , and fructooligosaccharides on bone growth and skeletal integrity of broiler chicks. *Poultry Sci.* 90, 2425–2432
- Littell R.C., Henry P.R., Lewis A.J., Ammerman C.B., 1997. Estimation of relative bioavailability of nutrients using SAS procedures. *J. Anim. Sci.* 75, 2672–2683

- Pillai P.B., O'Connor-Dennie T., Owens C.M., Emmert J.L., 2006. Efficacy of an *Escherichia coli* phytase in broilers fed adequate or reduced phosphorus diets and its effect on carcass characteristics. *Poultry Sci.* 85, 1737–1745
- Qian H., Komegay E.T., Denbow D.M., 1997. Utilization of phytate phosphorus and calcium as influenced by microbial phytase, cholecalciferol, and the calcium total phosphorus ratio in broiler diets. *Poultry Sci.* 76, 37–46
- Rao S.V.R., Raju M.V.L.N., Panda A.K., Shyam Sunder G., Sharma R.P., 2006. Effect of high concentrations of cholecalciferol on growth, bone mineralization and mineral retention in broiler chicks fed suboptimal concentrations of calcium and non-phytate phosphorus. *J. Appl. Poultry Res.* 131, 135–150
- Rao S.V.R., Raju M.V.L.N., Panda A.K., Shyam Sunder G., Sharma R.P., 2009. Performance and bone mineralisation in broiler chicks fed on diets with different concentrations of cholecalciferol at a constant ratio of calcium to non-phytate phosphorus. *Brit. Poultry Sci.* 50, 528–535
- SAS, 2001. SAS/STAT User's Guide. Release 8.2. Stat. Anal. Syst. SAS Institute Inc. Cary, NC
- Scheideler S.E., Ferket P.R., 2000. Phytase in broiler rations—effects on carcass yields and incidence of tibial dyschondroplasia. *J. Appl. Poultry Res.* 9, 468–475
- Snow J.L., Baker D.H., Parsons C.M., 2004. Phytase, citric acid, and 1 α -hydroxycholecalciferol improve phytate phosphorus utilization in chicks fed a corn-soybean meal diet. *Poultry Sci.* 83, 1187–1192
- Soares J.H. Jr., Swerdel M.R., Bossard E.H., 1978. Phosphorus availability. 1. The effect of chick age and vitamin D metabolites on the availability of phosphorus in defluorinated phosphate. *Poultry Sci.* 57, 1305–1312
- Whitehead C.C., McCormack H.A., McTeir L., Fleming R.H., 2004. High vitamin D₃ requirements in broilers for bone quality and prevention of tibial dyschondroplasia and interactions with dietary calcium, available phosphorus and vitamin A. *Brit. Poultry Sci.* 45, 425–436
- Yarger J.G., Saunders C.A., McNaughton J.L., Quarles C.L., Hollis B.W., Gray R.W., 1995. Comparison of dietary 25-hydroxycholecalciferol and cholecalciferol in broiler chickens. *Poultry Sci.* 74, 1159–1167